

# TANAGRA BRASILIA.



## CHARACTER GENERICUS.

*Rosstrum* conicum, acuminatum, emarginatum,  
basi subtrigonum, apice declive.

*Lin. Syst. Nat.* p. 314.

## CHARACTER SPECIFICUS, &c.

TANAGRA COCCINEA, alis caudaque nigris.

*Lin. Syst. Nat.* p. 314.

MERULA BRASILICA.

*Bell. av.* 319.

TIJEPIRANGA.

*Raj. syn.* p. 87. 13.

*Will. orn.* p. 184. 185.

CARDINALIS.

*Brieff. av.* 3. p. 42. n. 24. t. 3. f. 1.

---

Pulcherrima hæc avis, cujus magnitudinem naturale ostendit tabula, amat omnem Americam Australem, præcipue autem Brasiliam.







A. 1'

under published by F. P. Vaiden & Co. No. 11, Avenue Street.

THE  
BRASILIAN TANAGER.



GENERIC CHARACTER.

*Bill* conic, sharp-pointed, flogging at the tip and slightly emarginated; somewhat trigonal towards the base.

SPECIFIC CHARACTER, &c.

SCARLET TANAGER with black wings and tail.

Le CARDINAL.

*Briff. orn.* 3. p. 42. 24. pl. 3. f. 1.

BRASILIAN TANAGER.

*Lath. Syn.* 1. p. 215.

---

This highly elegant species is a native of South America, and is principally found in Brazil. The plate represents it in its natural size.